

Human Atherosclerotic Plaques Contain Gamma-Glutamyl Transpeptidase Enzyme Activity

Aldo Paolicchi, Michele Emdin, Erri Ghiozeni, Eugenio Ciancia, Claudio Passino, George Popoff and Alfonso Pompella

Circulation. 2004;109:1440

doi: 10.1161/01.CIR.0000120558.41356.E6

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2004 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/content/109/11/1440>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>


Subscriptions: Information about subscribing to *Circulation* is online at:
<http://circ.ahajournals.org/subscriptions/>

Human Atherosclerotic Plaques Contain Gamma-Glutamyl Transpeptidase Enzyme Activity

Aldo Paolicchi, MD; Michele Emdin, MD; Erri Ghiozeni, MD; Eugenio Ciancia, MD;
Claudio Passino, MD; George Popoff, MD; Alfonso Pompella, MD

During the last decade, growing evidence has shown that serum gamma-glutamyl transpeptidase (GGT) is an independent prognostic marker for cardiac death and reinfarction, both in unselected populations and in patients with coronary artery disease. Clinical and epidemiological evidence indicates that the prognostic value of GGT is largely independent of other risk factors for cardiovascular disease and alcohol consumption. The catalytic activity of GGT, which is present on the surface of cell membranes and in serum, is responsible for the extracellular

catabolism of the antioxidant glutathione. Cysteinyl glycine deriving from the hydrolysis of glutathione performed by GGT has been found to trigger iron-dependent production of reactive oxygen species as well as low-density lipoprotein oxidation in vitro. The localization of GGT within the coronary plaque (Figure) provides a pathological basis for the hypothesis of a direct participation of GGT in low-density lipoprotein oxidation within the plaque and in atherogenesis and coronary artery disease progression.



Histochemical and immunohistochemical demonstration of GGT enzyme activity within a frozen section of coronary atheroma from endoarterectomy in vivo. Histochemical reaction for GGT enzyme activity was performed by using the specific substrate gamma-glutamyl-4-methoxy-2-naphthylamide and the diazonium salt Fast Garnet GBC as a chromogen. A strong GGT activity (the red stain) is selectively present in correspondence of the core of the atheroma, while the fibrous cap stains negative (A, magnification 20 \times). The identification of the enzyme was confirmed immunohistochemically using a polyclonal antibody directed against the heavy chain of human GGT, which marked the same part of the atheroma (B and C, magnification 40 \times and 10 \times , respectively). The homogeneous immunostaining of oxidized lipid-containing foam cells by the antibody is evident at higher magnification (B). For the sake of confirming the localization of GGT activity, another section of the same specimen was immunostained with an antibody directed against CD 68+ cells (ie, macrophages), selectively identifying foam cells (D and E, magnification 10 \times and 40 \times , respectively).

From the Department of Experimental Pathology, University of Pisa, Pisa (A. Paolicchi, A. Pompella); the Institute of Clinical Physiology, National Research Council, Pisa (M.E., C.P.); Azienda Ospedaliera Pisana, Pisa (E.C.); and Clinica Villa Maria Beatrice, Firenze (E.G., G.P.), Italy.

Correspondence to Prof Aldo Paolicchi, Dipartimento di Patologia Sperimentale, sez. di Patologia Generale, Scuola Medica, via Roma, 55 I-56126, Pisa, Italy. E-mail paolicchi@biomed.unipi.it

The editor of Images in Cardiovascular Medicine is Hugh A. McAllister, Jr, MD, Chief, Department of Pathology, St Luke's Episcopal Hospital and Texas Heart Institute, and Clinical Professor of Pathology, University of Texas Medical School and Baylor College of Medicine.

Circulation encourages readers to submit cardiovascular images to the *Circulation* Editorial Office, St Luke's Episcopal Hospital/Texas Heart Institute, 6720 Bertner Ave, MC1-267, Houston, TX 77030.

(*Circulation*. 2004;109:1440.)

© 2004 American Heart Association, Inc.

Circulation is available at <http://www.circulationaha.org>

DOI: 10.1161/01.CIR.0000120558.41356.E6